



Independents Own

# ENCORE


POUR-ON LOUSICIDE FOR SHEEP

DON'T

LET LICE

SPOIL

THE SHOW



**Active Constituent:** 35 g/L Imidacloprid  
Also contains: 300 g/L N-METHYL-2-PYRROLIDONE (solvent)

**Claims:** For the control of neonicotinoid susceptible body lice *Bovicola ovis* on shorn sheep up to 7 days off shears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

**Container size:** 20L

**Applicator:** Apply product only with the 6-stripe applicator.

**Withholding Periods:** MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: DO NOT USE on female sheep which are producing, or may in the future produce, milk or milk products for human consumption.

WOOL: DO NOT USE less than 2 months before shearing or fibre collection.

**(ESI):** DO NOT slaughter for export less than 63 days after treatment.

**Other information on product:**

May be used in conjunction with worm control drenches and at the same time as vaccination.

Contains a blue dye to identify treated sheep. This dye is scourable from the wool.

Is safe for use on breeding rams.

### DOSAGE TABLE

BODYWEIGHT (KG)	DOSE (ML)	METHOD	APPLY AS BELOW
Below 6kg	8	Single Stripe	
6 – 8	12		
8.1 – 10	15		
10.1 – 12.5	20		
12.6 – 15	25		
15.1 – 20	30		
20.1 – 30	40	Double Stripe	
30.1 – 55	60		
55.1 – 80	80		
80.1 – 100	90		
Add 10mL for each 10kg above 100kg			

FOR LABEL, SDS AND ALL TECHNICAL ENQUIRIES, PLEASE GO TO:  
[independentsown.com.au](http://independentsown.com.au) E: [info@airr.com.au](mailto:info@airr.com.au) P: (03) 5821 0922

APVMA Approval No.: 84054/118626