

85mm

185mm

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ENVIRONMENTAL PROTECTION

Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of container and outer packaging by wrapping with paper and putting in garbage.

STORAGE

Store below 30°C (room temperature). Store bottle in carton to protect from light.

APVMA Approval No.: 90419/131992

Harmful if contact with skin. Causes serious eye irritation. May damage fertility or the unborn child.

May cause harm to breast-fed children.

Do not handle until all safety precautions have been read and understood.

Avoid contact during pregnancy or while nursing. Wash contacted areas thoroughly after handling.

If exposed or concerned or if eye irritation persists: Get medical advice.

Store locked up and in a well-ventilated place. Keep cool.



**UN 2902
PESTICIDE,
LIQUID,
TOXIC,
N.O.S., PG III**

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



MECTEC PLUS

(IVERMECTIN PLUS CLORSULON)

For the treatment and control of ivermectin and clorsulon sensitive strains of internal and external parasites of cattle, including adult liver flukes.



ACTIVE CONSTITUENTS:

100 mg/mL CLORSULON, 10 mg/mL IVERMECTIN

Made for AIRR 74-76 Drummond Rd, Shepparton VIC 3630
P 03 5820 8400 independentsown.com.au

CONTENTS
500
ML



iO/10
GUARANTEE

PROUDLY

OWNED & MADE

Read the carton for full instructions.

DIRECTIONS FOR USE

Restraints:

Re-treatment interval: **DO NOT** re-treat animals for 28 days after last treatment.

Dosage and Administration:

To be given by subcutaneous injection only at the recommended dose of 1 mL per 50 kg liveweight (equivalent to 0.2 mg ivermectin per kg and 2 mg clorsulon per kg liveweight). Refer to carton for dosage table.

Inject under loose skin high on the neck, behind the ear in an area away from the more valuable carcass muscle. Loose skin on the neck in front of the shoulder is also a suitable site. This product is not to be used intravenously or intramuscularly.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: Zero (0) days. There is a NIL milk withholding period for lactating dairy cattle.

TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI) **DO NOT USE** less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from AIRR on (03) 5820 8400 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes, skin and clothing.

BATCH:

EXPIRY:

31952F-1